

# ECOTOXICOLOGY OF MULCH FILMS



For optimal scanning, please enlarge poster to 125%.  
antonia.weltmeyer@rwth-aachen.de

Antonia Weltmeyer<sup>1</sup>, Christoph Kämpfer<sup>1</sup>, Thomas-Benjamin Seiler<sup>1</sup>, Hennert Hollert<sup>1,2</sup>, Martina Roß-Nickoll<sup>1</sup>

<sup>1</sup> RWTH Aachen University, <sup>2</sup> Goethe University Frankfurt

## BACKGROUND

- Mulch films are applied worldwide due to their **positive effects** (e.g. humidity, temperature, weed growth limitation) on agricultural farming
- Commonly applied polymers are polyethylene (PE) and the bio-based poly lactate acid (PLA) with addition of multiple, often **unknown additives**
- Despite the environmentally friendly seeming labelling of bio-based and biodegradable, PLA can be even **more toxic** than petroleum based plastics<sup>2</sup> and only degraded under very harsh conditions<sup>3</sup>
- Degradation and later accumulation of mulch films in soil is facilitated by their low thickness due to rising removal difficulty and generally high disposal costs<sup>4</sup>
- Plastic mulching is a **relevant source of microplastics** (MP) in the terrestrial environment with up to 324.5 kg/ha<sup>5</sup> which lies in the same magnitude of earthworm mortality<sup>6</sup>
- Impact of accumulation in soil is yet insufficiently determined and requires further investigation<sup>7</sup>

The main applications, greenhouse and mulch film, account for almost **75%** of the total market, with more than one million tons being produced for each.<sup>1</sup>

Plasticulture market growth is **5% annually** depending on region and application.<sup>1</sup>



We aim to investigate the effects of mulch film microplastics and their constituents on the living environment with focus on soil organisms.

Furthermore, we want to investigate the influence of sorbed pesticides on the toxicity of mulch film fragments.



This work is part of the iMulch project funded by the European Fond for regional development (EFRE).

iMulch – An investigation of the influence of polymers on a terrestrial ecosystem using the example of mulch films used in agriculture

## OBJECTIVE

- Assessment of toxicity mechanisms of mulch film constituents and potential synergistic effects
- Determination of MP toxicity on representative soil organisms
- Determination of the interaction of mulch film fragments and selected pesticides
  - Influence of mulch films on pesticide toxicity
  - Availability of pesticides in plastic spiked soil
- Determination of the baseline toxicity of pesticide active substances and products



## WHERE DO WE GO FROM HERE?

- In vitro**
- Establishment of beta-Naphtoflavone as micro EROD induction standard
  - Investigation of sorption behaviour in microtiter plates for extracts and pesticides
- In vivo**
- Comparison of effects of new and aged mulch film
  - Deduction for further testing with *Eisenia fetida* in chronic exposure scenarios (OECD 222, in soil) and decision on biomarkers
  - Comparison between sensitivity of *Eisenia fetida* and *Lumbricus terrestris*



Other project partners are also presenting at the SETAC 2020: Check out poster 2.08P.11 for information on biodegradability and poster 4.14P.1 for more information on the iMulch project and first results.

## METHODS



### Extraction

- 500mg polymer cut into 1x2cm
- Worst case scenario**
- Per solvent 2x30min UAE
- Solvents PE: acetonitrile/dichlormethane  
Solvents PLA: acetonitrile / isopropanol
- Realistic scenario**
- Shaking extraction: 24h, Millipore Water, 20±2°C, 100rpm
- Solid phase extraction
  - Solvent: ethyl acetate, methanol
- Analysis with GC-MS

### Extracts & pesticide screening

- Extract testing
  - In vitro screening for dioxin-like activity, genotoxicity and endocrine/androgene activity
  - Application of glass-coated microplates to prohibit sorption
  - Determination of acute toxicity to earthworm *Eisenia fetida* (OECD 207, filter paper test)
- Pesticide testing
  - In vitro assays to narrow down substances
  - Active substances and products in comparison

### Microplastic in soil

- Reference soil 01-A to approximate to natural conditions
- Scenarios**
- Pesticide loaded MP vs. MP + pesticide
- Exposure monitoring (soil, earthworm)
  - Acute toxicity on *E. fetida* (OECD 207)
  - OECD 232: realistic and worst case MP concentrations on *F. candida* reproduction
- Details on MP production:

## PRELIMINARY RESULTS

✓      ✗      ?

- In vitro pesticide screening: positive, negative, unclear

	Endocrine activity	Genotoxicity	Dioxin-like activity
Atrazine	✓	✓	?
Tebuconazole	✓	✗	✓
Quinoxifen	✓	?	✗

- 2,3,7,8-TCDD not usable in assay with glass coated plates (Fig. 1)
- OECD 207: Filterpaper test with EC50 [mg/mL]
  - Tebuconazole (49±2), solvent extracts (no toxicity), water extracts (no toxicity)
- OECD 207: Acute toxicity test with active substance and EC50 [mg/mL]
  - Folicur (Tebuconazole, 116±27), Biscaya (Thiachloprid, <50)




Fig. 1 EROD activity for 2,3,7,8-TCDD in plastic plates (blue, n=5) and glass plates (n=4)

## REFERENCES

- BASF, 2016. Plastic additives for Agricultural Plastics, available at: [https://agriculture.basf.com/global/assets/en/Crop%20Protection/use-areas/Agriculture\\_2016\\_Final.pdf](https://agriculture.basf.com/global/assets/en/Crop%20Protection/use-areas/Agriculture_2016_Final.pdf) (last access: 15.04.2020)
  - Zimmermann, L., Dierkes, G., Ternes, T. A., Völker, C. and Wagner, M. Benchmarking the in Vitro Toxicity and Chemical Composition of Plastic Consumer Products. Environ. Sci. Technol. 53, 19, 11467-11477 (2019). <https://doi.org/10.1021/acs.est.9b02293>
  - Haider, T. P., Völker, C., Kramm, J., Landfester, K. and Wurm, F. R. Plastics of the Future? The Impact of Biodegradable Polymers on the Environment and on Society Angew. Chem. Int. Ed. 58, 50-62 (2019). <https://doi.org/10.1002/anie.201805766>
  - OWS, 2017. Expert Statement: (Bio)Degradable Mulching Films, available at: <http://www.ows.be/wp-content/uploads/2017/08/Expert-statement-mulching-films.pdf> (last access: 15.04.2020)
  - Huang, Y., Liu, Q., Jia, W., Yan, C. and Wang, J. Agricultural plastic mulching as a source of microplastics in the terrestrial environment. Environ. Poll. 260, 114096, (2020). <https://doi.org/10.1016/j.envpol.2020.114096>
  - Cao, D., Wang, X., Luo, X., Liu, G. and Zheng, H. Effects of polystyrene microplastics on the fitness of earthworms in an agricultural soil. IOP Conf. Ser.: Earth Environ. Sci. 61 012148 (2017). <http://dx.doi.org/10.1088/1755-1315/61/1/012148T1>
  - Zhu, F., Zhu, C. and Wang, C. et al. Occurrence and Ecological Impacts of Microplastics in Soil Systems: A Review. Bull Environ Contam. Toxicol. 102, 741-749 (2019). <https://doi.org/10.1007/s00128-019-02623-z>
- Picture: Source: [https://www.basf.com/global/en/products/plastics-rubber/fairs/BASFatK2016/must\\_sees/ecovio\\_m.html](https://www.basf.com/global/en/products/plastics-rubber/fairs/BASFatK2016/must_sees/ecovio_m.html) (last access: 19.04.2020)